

ABOUT THE AUTHOR

Jennifer Tran, MD

Dr. Jennifer Tran is a board-certified dermatologist in Canada and the U.S. with expertise in medical, surgical, and cosmetic dermatology. Dr. Tran completed her undergraduate training in Health Sciences at McMaster University. She then completed her medical degree and subsequently her dermatology residency at the University of Toronto, where she served as chief resident during her final year of training. She is currently working at several community clinics and runs an academic practice at Sunnybrook Health Sciences Centre.



AN OVERVIEW OF LICHEN SCLEROSUS

INTRODUCTION

Lichen sclerosus (LS) is a chronic, inflammatory disease that typically affects the genital skin, and less frequently on other skin sites. It is characterized by white, shiny, scar-like atrophy. Symptoms include pruritus, burning, progressive atrophy, and in severe cases, loss of normal genital architecture, and functional impairment.

Synonyms: kraurosis vulvae, balanitis xerotic obliterans (in males), lichen sclerosus et atrophicus

EPIDEMIOLOGY

The exact incidence and prevalence of lichen sclerosus are not known due to a lack of population-based data in the literature. One previous review of a general gynecology practice estimated the prevalence of vulvar LS to be around 1.7%¹ and a Dutch cohort study estimated the incidence rate of lichen sclerosus to range from 7.4 to 14.6 per 100,000 woman-years between 1991 and 2011.² Peak incidence occurs in postmenopausal women, with a second peak in prepubertal girls, followed by men in their fourth decade.³

ETIOLOGY/PATHOGENESIS

The cause of LS remains unknown. Several HLA haplotypes have been associated with the disease and identified in the peer-reviewed literature.

(HLA-DQ3, HLA-DQ7, HLA-DRB1*12, HLA-DRB1*0301/04/DQB1*0201/02/03, among others).⁴⁻⁶

Autoantibodies against extracellular matrix protein 1 (ECM-1) have also been identified in LS patients. Indeed, autoimmunity is believed to play a primary role in the disease, and several autoimmune comorbid conditions have been associated with LS, including autoimmune thyroiditis, alopecia areata, vitiligo, pernicious anemia, celiac disease, and morphea.^{7,8} *Borrelia burgdorferi sensu lato* species has also been implicated in the pathogenesis of lichen sclerosus, with one study detecting *Borrelia* species in 63% (38 of 60 cases) of lichen

sclerosus tissue specimens.⁹ Other triggers, such as influenza vaccination¹⁰, surgery¹¹, and genital piercing¹² have been described as well in the form of single case reports.

CLINICAL FEATURES

LS can affect both the genital and extragenital skin. The classic morphology is characterized by ivory white, porcelain-like, sclerotic, atrophic papules and plaques. These lesions may also have telangiectasias, follicular plugging, erosions, purpura, and haemorrhagic bullae. In the anogenital region in women, a “figure-of-eight” configuration has been described, with circumferential lesions involving the vulva, perineum and anus. In advanced cases, the normal architecture of the labia minora, labia majora, and clitoral hood may be resorbed, resulting in narrowing of the vaginal introitus. In men, lesions may similarly present as sclerotic, shiny, white lesions, often on the glans penis and inner aspect of the foreskin. Constriction, phimosis, paraphimosis, and recurrent balanitis are complications of LS in men. In advanced stages, patients may also have urinary obstruction. Patients with LS, both men and women, often have significant dyspareunia, pruritus, dysuria, and general soreness and discomfort.

Because of the chronic inflammatory state of LS, there is a risk of malignancy in longstanding lesions. The 10-year vulvar squamous cell carcinoma incidence in women with lichen sclerosus was associated with concurrent vulvar intraepithelial neoplasia and age at time of lichen sclerosus diagnosis (5.9% in women of ≥ 70 years, 3% in women between 50 and 70 years, and 1.8% in women < 50 years).²

DIFFERENTIAL DIAGNOSIS

Lichen sclerosus must be delineated from a variety of other diseases of the anogenital region, including morphea, erosive lichen planus, and graft vs host disease. In men, Erythroplasia of Queyrat, balanitis, and extramammary Paget's must all be considered. In young children, it is important to rule out sexual abuse.^{3,13,14}

PATHOLOGY

Biopsies of lichen sclerosus will typically show a thinned epidermis and a lichenoid interface dermatitis characterized by vacuolar degeneration of the basal layer and a band-like lymphocytic infiltrate.¹³⁻¹⁶ Macrophages and mast cells can also sometimes be seen in the infiltrate. Follicular plugging may be seen and rete ridges may be flattened. Below this, in the dermal level, lesions may demonstrate homogenized dermal collagen and edema. The key differentiating feature between lichen sclerosus and morphea is the loss of elastic fibers, which is seen in LS but not morphea.¹⁴⁻¹⁶

TREATMENT

The current standard of care for LS patients is a high potency topical steroid such as clobetasol propionate 0.05% daily or alternating with a non-steroidal topical or rest period.^{3,14,17} For recalcitrant cases, or for patients unable to apply topical steroids, intralesional steroid injections can be helpful.¹⁷ The most commonly used non-steroidal alternatives are tacrolimus and pimecrolimus. One study of sixteen patients (10 with anogenital and six with extragenital localization) were treated with topical tacrolimus ointment twice daily. Results demonstrated improvement in a majority of genital LS patients¹⁸, but none in patients with extragenital LS.

No adverse effects were observed in this study. Due to a black box warning regarding lymphoma risk associated with tacrolimus and the risk of squamous cell carcinoma (SCC) arising in LS, clinicians should consider counselling patients about this theoretical risk. However, no studies have shown development of SCC as a result of topical calcineurin inhibitors in patients with LS.

Although topical estrogen and progesterone preparations have been used for post-menopausal dryness and dyspareunia, there is no evidence to support the use of such preparations in LS.¹⁷ Topical testosterone is also not recommended and has been shown to be inferior to clobetasol.¹⁷ Furthermore, studies have demonstrated unacceptable adverse effects of topical testosterone including clitoral hypertrophy, hirsutism, acne, and menstrual abnormalities.^{17,19,20}

Phototherapy has long been considered in the treatment of LS. The most evidence exists for the use of PUVA/UVA1, but the data is limited to a few case series, and patients should be warned about risk of carcinoma with PUVA.

The use of systemic treatments such as prednisone, cyclosporine, and methotrexate have been elucidated through case reports in the literature. The largest study of methotrexate was a retrospective review of 28 patients who had previously failed topical therapies. The authors found improvement in 75% of patients who received weekly doses of methotrexate ranging from 2.5 mg to 17.5 mg (median = 10 mg), however several patients had to discontinue methotrexate due to side effects.²¹

There are conflicting case reports involving the use of hydroxychloroquine and thus a recent review of treatment strategies for LS¹⁷ did not recommend this medication.

There is some evidence for the use of systemic retinoids in LS. One randomized controlled trial²² found a significantly higher number of responders in patients receiving 35 mg of acitretin daily (14 of 22 patients), compared with the placebo group (6 of 24 patients). Another randomized, double-blind, placebo controlled trial done in males with LS²³ also found some positive effect of acitretin. In this study of 49 completers who were eligible for statistical analysis, complete response was achieved by 36.4% (12 of 33) of the acitretin group vs 6.3% (1 of 16) of the controls, while 36.4% (12 of 33) vs 12.5% (2 of 16) achieved partial resolution, respectively.

A potential hypothesis for the treatment of LS centers on the use of systemic antibiotics, which links back to the theory that *Borrelia* species may trigger LS. One retrospective study of 15 men and women with steroid-resistant LS were treated with one of: intramuscular penicillin and oral penicillin, intramuscular cephalosporin, oral penicillin, or oral cephalosporin.²⁴ The authors noted that all patients showed significant response within a few weeks, particularly those who had received intramuscular antibiotics, with significant reduction in pain, pruritus, and burning. Though not a first-line treatment approach, clinicians may consider the use of intramuscular ceftriaxone 1 gram every two weeks for three doses,

then once a month on a PRN basis; or intramuscular penicillin G benzathine suspension dosed at 2.4 million units every 2 weeks for three doses, then once a month on a PRN basis in resistant LS cases.¹⁷

Newer, investigative treatments include the use of fractional carbon dioxide laser (i.e. MonaLisa Touch®) which is currently being investigated at Sunnybrook Health Sciences Centre. The published data on this treatment modality is limited, with mostly case reports reported in the literature.²⁵⁻²⁸

The largest published study of forty women with LS who failed clobetasol propionate demonstrated improvement in vulvar itching, dryness, sensitivity during intercourse and dyspareunia. Itching relief was noted after only one treatment in several patients.²⁹ No systemic or local adverse reactions were reported in this study. Fractional CO2 laser has also been studied in males, with improvement noted in DLQI scores, symptoms of LS, sexual function, and no relapse at the 6-month follow-up mark.³⁰ This study also reported that the fractional laser was well-tolerated, with no significant adverse effects. Some patients experienced edema, short duration of burning, and erythema, all of which resolved within hours to days.

Surgical interventions have also been studied in LS. In men, surgical intervention such as circumcision can often be curative in mild-to-moderate cases.^{13,17} In women with advanced LS, perineotomy and lysis of adhesions can provide symptomatic relief.¹⁷ Platelet-rich plasma has also been studied in a non-randomized setting, with excellent response rates in subjects ranging from 62%-100%.³¹⁻³⁵

SUMMARY

Lichen sclerosus is a chronic, inflammatory disease that primarily affects the anogenital area in both men and women. Treatment is essential to prevent progression of the disease, which can lead to permanent deformity and decreased quality of life. The mainstay of treatment includes potent topical steroids and in some cases, surgery can be curative. In cases unresponsive to topical steroids, systemic treatments such as systemic retinoids, methotrexate, and antibiotics may be utilized. Fractional laser and platelet-rich plasma are promising new treatments and may be considered in recalcitrant cases.

1. Goldstein AT, Marinoff SC, Christopher K, Srodon M. Prevalence of vulvar lichen sclerosis in a general gynecology practice. *J Reprod Med.* 2005;50(7):477-480.
2. Bleeker MC, Visser PJ, Overbeek LI, van Beurden M, Berkhof J. Lichen Sclerosus: Incidence and Risk of Vulvar Squamous Cell Carcinoma. *Cancer Epidemiol Biomarkers Prev.* 2016;25(8):1224-1230. doi:10.1158/1055-9965.EPI-16-0019
3. Fistarol SK, Itin PH. Diagnosis and treatment of lichen sclerosis: an update. *Am J Clin Dermatol.* 2013;14(1):27-47. doi:10.1007/s40257-012-0006-4
4. Gao XH, Barnardo MC, Winsey S, et al. The association between HLA DR, DQ antigens, and vulval lichen sclerosis in the UK: HLA DRB1*12 and its associated DRB1*12/DQB1*03:01/04/010 haplotype confers susceptibility to vulval lichen sclerosis, and HLA DRB1*03:01/04 and its associated DRB1*03:01/04/DQB1*02:01/03 haplotype protects from vulval lichen sclerosis. *J Invest Dermatol.* 2005;125(5):895-899. doi:10.1111/j.0022-202X.2005.23905.x
5. Marren P, Yell J, Charnock FM, Bunce M, Welsh K, Wojnarowska F. The association between lichen sclerosis and antigens of the HLA system. *Br J Dermatol.* 1995;132(2):197-203. doi:10.1111/j.1365-2133.1995.tb05013.x
6. Powell J, Wojnarowska F, Winsey S, Marren P, Welsh K. Lichen sclerosis premenarche: autoimmunity and immunogenetics. *Br J Dermatol.* 2000;142(3):481-484. doi:10.1046/j.1365-2133.2000.03360.x
7. Meyrick Thomas RH, Ridley CM, McGibbon DH, Black MM. Lichen sclerosis et atrophicus and autoimmunity--a study of 350 women. *Br J Dermatol.* 1988 Jan;118(1):41-6. doi: 10.1111/j.1365-2133.1988.tb01748.x. Erratum in: *Br J Dermatol* 1988 May;118(5):736. PMID: 3342175.
8. Cooper SM, Ali I, Baldo M, Wojnarowska F. The association of lichen sclerosis and erosive lichen planus of the vulva with autoimmune disease: a case-control study. *Arch Dermatol.* 2008 Nov;144(11):1432-5. doi: 10.1001/archderm.144.11.1432. PMID: 19015417.
9. Eisendle K, Grabner T, Kutzner H, Zelger B. Possible role of *Borrelia burgdorferi* sensu lato infection in lichen sclerosis. *Arch Dermatol.* 2008;144(5):591-598. doi:10.1001/archderm.144.5.591
10. Requena López S, Hidalgo García Y, Gómez Díez S, Vivanco Allende B. Morphea and Extragenital Lichen Sclerosus et Atrophicus After Influenza Vaccination. Morfea y liquen escleroatrófico extragenital generalizados tras vacuna antigripal. *Actas Dermosifiliogr.* 2018;109(1):86-88. doi:10.1016/j.ad.2017.05.016
11. Pass CJ. An unusual variant of lichen sclerosis et atrophicus: delayed appearance in a surgical scar. *Cutis.* 1984;33(4):405-408.
12. De Giorgi V, Scarfi F, Silvestri F, et al. Genital piercing: A warning for the risk of vulvar lichen sclerosis. *Dermatol Ther.* 2021;34(1):e14703. doi:10.1111/dth.14703
13. Fergus KB, Lee AW, Baradaran N, et al. Pathophysiology, Clinical Manifestations, and Treatment of Lichen Sclerosus: A Systematic Review. *Urology.* 2020;135:11-19. doi:10.1016/j.urology.2019.09.034
14. Bolognia JL, Schaffer JV, Cerroni L. *Dermatology.* 4th edition. Elsevier; 2018
15. Patterson, J. *Weedon's Skin Pathology.* 4th Edition. Elsevier; 2014
16. Rapini, R. *Practical Dermatopathology.* Elsevier, 2012.
17. Kirtschig G, Becker K, Günthert A, et al. Evidence-based (S3) Guideline on (anogenital) Lichen sclerosis. *J Eur Acad Dermatol Venereol.* 2015;29(10):e1-e43. doi:10.1111/jdv.13136
18. Kim GW, Park HJ, Kim HS, et al. Topical tacrolimus ointment for the treatment of lichen sclerosis, comparing genital and extragenital involvement. *J Dermatol.* 2012;39(2):145-150. doi:10.1111/j.1346-8138.2011.01384.x
19. Cattaneo A, Carli P, De Marco A, et al. Testosterone maintenance therapy. Effects on vulvar lichen sclerosis treated with clobetasol propionate. *J Reprod Med.* 1996;41(2):99-102.
20. Bornstein J, Heifetz S, Kellner Y, Stolar Z, Abramovici H. Clobetasol dipropionate 0.05% versus testosterone propionate 2% topical application for severe vulvar lichen sclerosis. *Am J Obstet Gynecol.* 1998;178(1 Pt 1):80-84. doi:10.1016/s0002-9378(98)70631-3
21. Cuellar-Barboza A, Bashyam AM, Ghamrawi RI, Aickara D, Feldman SR, Pichardo RO. Methotrexate for the treatment of recalcitrant genital and extragenital lichen sclerosis: A retrospective series. *Dermatol Ther.* 2020;33(4):e13473. doi:10.1111/dth.13473
22. Bousema MT, Romppanen U, Geiger JM, et al. Acitretin in the treatment of severe lichen sclerosis et atrophicus of the vulva: a double-blind, placebo-controlled study. *J Am Acad Dermatol.* 1994;30(2 Pt 1):225-231. doi:10.1016/s0190-9622(94)70021-4
23. Ioannides D, Lazaridou E, Apalla Z, Sotiriou E, Gregoriou S, Rigopoulos D. Acitretin for severe lichen sclerosis of male genitalia: a randomized, placebo controlled study. *J Urol.* 2010;183(4):1395-1399. doi:10.1016/j.juro.2009.12.057
24. Shelley WB, Shelley ED, Amurao CV. Treatment of lichen sclerosis with antibiotics. *Int J Dermatol.* 2006;45(9):1104-1106. doi:10.1111/j.1365-4632.2006.02978.x
25. Gardner AN, Aschkenazi SO. The short-term efficacy and safety of fractional CO2 laser therapy for vulvovaginal symptoms in menopause, breast cancer, and lichen sclerosis [published online ahead of print, 2021 Jan 4]. *Menopause.* 2021; Publish Ahead of Print:10.1097/GME.0000000000001727. doi:10.1097/GME.0000000000001727
26. Mendieta-Eckert M, Torrontegui Bilbao J, Zabalza Estévez I, Landa Gundin N. Treatment of Vulvar Lichen Sclerosus et Atrophicus With Fractional Carbon Dioxide Laser Therapy: A Report of 4 Cases. *Tratamiento de liquen escleroso y atrófico vulvar con láser de dióxido de carbono fraccionado. Presentación de cuatro casos. Actas Dermosifiliogr.* 2021;112(1):85-88. doi:10.1016/j.ad.2019.03.021
27. Mendieta-Eckert M, Ocerin-Guerra I, Landa-Gundin N. Lichen sclerosis et atrophicus in a surgical scar treated with fractional laser. *J Cosmet Laser Ther.* 2017;19(2):106-108. doi:10.1080/14764172.2016.1262955
28. Lee A, Lim A, Fischer G. Fractional carbon dioxide laser in recalcitrant vulvar lichen sclerosis. *Australas J Dermatol.* 2016;57(1):39-43. doi:10.1111/ajd.12305
29. Pagano, Tiziana, et al. "Effect of rescue fractional microablative CO2 laser on symptoms and sexual dysfunction in women affected by vulvar lichen sclerosis resistant to long-term use of topic corticosteroid: a prospective longitudinal study." *Menopause* 27.4 (2020): 418-422.
30. Ferrara F, Messori S, Abbenante D, Patrizi A, Bardazzi F. Fractional CO2 laser therapy of lichen sclerosis in males: a new therapeutic opportunity? [published online ahead of print, 2020 Jul 20]. *J Dermatolog Treat.* 2020;1-5. doi:10.1080/09546634.2020.1793886
31. Tedesco M, Pranteda G, Chichierchia G, et al. The use of PRP (platelet-rich plasma) in patients affected by genital lichen sclerosis: clinical analysis and results. *J Eur Acad Dermatol Venereol.* 2019;33(2):e58-e59. doi:10.1111/jdv.15190
32. Behnia-Willison F, Pour NR, Mohamadi B, et al. Use of Platelet-rich Plasma for Vulvovaginal Autoimmune Conditions Like Lichen Sclerosus. *Plast Reconstr Surg Glob Open.* 2016;4(11):e1124. Published 2016 Nov 23. doi:10.1097/GOX.0000000000001124
33. Casabona F, Gambelli I, Casabona F, Santi P, Santori G, Baldelli I. Autologous platelet-rich plasma (PRP) in chronic penile lichen sclerosis: the impact on tissue repair and patient quality of life. *Int Urol Nephrol.* 2017;49(4):573-580. doi:10.1007/s11255-017-1523-0
34. Navarrete J, Echarte L, Sujánov A, et al. Platelet-rich plasma for male genital lichen sclerosis resistant to conventional therapy: First prospective study. *Dermatol Ther.* 2020;33(6):e14032. doi:10.1111/dth.14032
35. Tedesco M, Garelli V, Bellei B, et al. Platelet-rich plasma for genital lichen sclerosis: analysis and results of 94 patients. Are there gender-related differences in symptoms and therapeutic response to PRP? [published online ahead of print, 2020 Dec 6]. *J Dermatolog Treat.* 2020;1-5. doi:10.1080/09546634.2020.1854650